

Large localizations of finite simple groups

RÜDIGER GÖBEL, JOSÉ L. RODRÍGUEZ AND SAHARON SHELAH *

February 1, 2008

Abstract

A group homomorphism $\eta : H \rightarrow G$ is called a *localization* of H if every homomorphism $\varphi : H \rightarrow G$ can be ‘extended uniquely’ to a homomorphism $\Phi : G \rightarrow G$ in the sense that $\Phi\eta = \varphi$.

Libman showed that a localization of a finite group need not be finite. This is exemplified by a well-known representation $A_n \rightarrow SO_{n-1}(\mathbb{R})$ of the alternating group A_n , which turns out to be a localization for n even and $n \geq 10$. Dror Farjoun asked if there is any upper bound in cardinality for localizations of A_n . In this paper we answer this question and prove, under the generalized continuum hypothesis, that every non abelian finite simple group H , has arbitrarily large localizations. This shows that there is a proper class of distinct homotopy types which are localizations of a given Eilenberg–Mac Lane space $K(H, 1)$ for any non abelian finite simple group H .

0 Introduction

One of the current problems in localization of groups is to decide what algebraic properties of H can be transferred to G by a localization $\eta : H \rightarrow G$. Recall that $\eta : H \rightarrow G$ is a *localization* if every homomorphism $\varphi : H \rightarrow G$ in the diagram

$$\begin{array}{ccc} H & \xrightarrow{\eta} & G \\ \varphi \downarrow & \swarrow \Phi & \\ G & & \end{array} \quad (0.1)$$

*The first and the third authors are supported by the project No. G 0545-173, 06/97 of the German-Israeli Foundation for Scientific Research & Development. The second author is supported by DGES grant PB97-0202 and DAAD. GbSh 701 in Shelah’s list of publications.

can be extended to a *unique* homomorphism $\Phi : G \rightarrow G$ such that $\Phi\eta = \varphi$. This in other words says that $G \cong L_\eta H$ where L_η is the localization functor with respect to η ; see e.g. [2].

For example, the properties of being an abelian group or a commutative ring with 1 are preserved. Casacuberta, Rodríguez and Tai [3] have found consequences of these facts for homotopical localizations of abelian Eilenberg–Mac Lane spaces (see also Casacuberta [2], Dror Farjoun [5]).

A further step leads to nilpotent groups. Dwyer and Dror Farjoun showed that every localization of a nilpotent group of class 2 is nilpotent of class 2 or less. A proof is given by Libman in [11]. However, it is unknown if nilpotent groups are preserved under localizations in general.

Another interesting problem is to find an upper bound for the cardinalities of the localizations of a fixed group H . It is easy to see that, if H is finite abelian, then every localization $\eta : H \rightarrow G$ is an epimorphism, hence $|G| \leq |H|$. More generally, Libman has shown in [11] that if H is torsion abelian then $|G| \leq |H|^{\aleph_0}$. However, if H is not torsion, this can fail. Indeed, the localizations of $\mathbb{Z}$ are precisely the E -rings, and we know by Dugas, Mader and Vinsonhaler [7] that there exist E -rings of arbitrarily large cardinality. Strüngmann sharpened this result in [17] for almost free E -rings.

The first example observing that a localization of a finite group need not be finite is due to Libman [12]. He showed that the alternating group A_n has a $(n-1)$ -dimensional irreducible representation $\varphi : A_n \rightarrow SO_{n-1}(\mathbb{R})$ which is a localization for any even $n \geq 10$. In the proof he uses that $O_{n-1}(\mathbb{R})$ is complete, $SO_{n-1}(\mathbb{R})$ is simple, and the fact that all automorphisms of $SO_{n-1}(\mathbb{R})$ are conjugation by some element in $O_{n-1}(\mathbb{R})$. This also motivates our Definition 1. Thus, Dror Farjoun asked about the existence of an upper bound for the cardinality of localizations of A_n . We give an answer to this question in Corollary 3, which is a direct consequence of our Main Theorem and Proposition 2.

In fact our result also holds for many other groups which we will call *suitable* groups, see Definition 1. However, we shall assume the generalized continuum hypothesis. GCH will be needed to apply a new combinatorial principle Dl_λ due to Hart, Laflamme and Shelah [9]. If λ is a successor of a regular cardinal, then under GCH this principle holds and is a general tool which allows us to manufacture objects of size λ^+ from approximations of size less than λ .

The group theoretical techniques derive from standard combinatorial group theory and can be found in the book by Lyndon and Schupp [13]. In particular, we will use small cancellation theory as in [16], where Shelah solved Kurosh's problem about the existence of Jonsson groups.

Main Theorem (*Assume ZFC + GCH*) *Let λ be the successor of an uncountable, regular cardinal and let λ^+ be its successor cardinal. Then any suitable group H whose automorphism group $\text{Aut}(H)$ has size $< \lambda$ is a subgroup of a group $\mathcal{G}$ of size λ^+ with the following properties:*

- (a) $\mathcal{G}$ is generated by copies of H .
- (b) Any monomorphism $\phi : H \rightarrow \mathcal{G}$ is the restriction of an inner automorphism of $\mathcal{G}$.
- (c) H has trivial centralizer in $\mathcal{G}$, i.e. if $x \in \mathcal{G}$ commutes with all elements of H , then $x = 1$.
- (d) $\mathcal{G}$ is complete, i.e. all automorphisms of $\mathcal{G}$ are inner automorphisms and $\mathcal{G}$ has trivial centre $Z(\mathcal{G})$.
- (e) $\mathcal{G}$ is co-hopfian, i.e. any monomorphism $\mathcal{G} \rightarrow \mathcal{G}$ is an automorphism.
- (f) $\mathcal{G}$ is simple.

Definition 1 Let H be any group with trivial center and view $H \leq \text{Aut}(H)$ as inner automorphisms of H . Then H is called *suitable* if the following conditions hold:

1. H is torsion, i.e. all elements have finite order.
2. If $H_1 \leq \text{Aut}(H)$ and $H_1 \cong H$ then $H_1 = H$.
3. H is complete in $\text{Aut}(H)$, i.e. any automorphism of H extends to an inner automorphism of $\text{Aut}(H)$.

We want to show that all non abelian finite simple groups are suitable. It only remains to show conditions 2 and 3 in the above definition. As a consequence of the classification of finite simple groups the Schreier conjecture holds for all finite groups H , hence the outer automorphism group $\text{Out}(H)$ is always soluble. If H is also simple and non abelian then H is a characteristic subgroup of $\text{Aut}(H)$. Hence any copy of H in

$\text{Aut}(H)$ is H and any automorphism of H is induced by an automorphism of $\text{Aut}(H)$. However, $\text{Aut}(H)$ is complete by Burnside, see [14, p. 399]. This is the first part of the following:

Proposition 2 (a) *All non abelian finite simple groups are suitable.*

(b) *The non abelian, finite, simple and complete groups are precisely the following sporadic groups: $M_{11}, M_{23}, M_{24}, Co_3, Co_2, Co_1, Fi_{23}, Th, B, M, J_1, Ly, Ru, J_4$.*

Part (b) follows by inspection of the list of finite simple groups. It is interesting to know when $H = \text{Aut}(H)$ because in this case our proof becomes visibly simpler.

We finally obtain an answer to Dror Farjoun's question.

Corollary 3 (Assume ZFC + GCH.) *Any suitable simple group has localizations of arbitrarily large cardinality.*

The localization $A_n \rightarrow SO_{n-1}(\mathbb{R})$ for any even $n \geq 10$ induces a map between Eilenberg–Mac Lane spaces $K(A_n, 1) \rightarrow K(SO_{n-1}(\mathbb{R}), 1)$ which turns out to be a localization in the homotopy category; [12]. This is the first example of a space with a finite fundamental group which admits localizations with an infinite fundamental group. Corollary 3 yields then the following extension:

Corollary 4 (Assume ZFC + GCH.) *Let H be a suitable simple group. Then $K(H, 1)$ has localizations with arbitrarily large fundamental group.*

Other constructions in homotopy theory based on large-cardinal principles were described by Casacuberta, Scevenels and Smith in [4].

Acknowledgements. We would like to thank Emmanuel Dror Farjoun for explaining the above problem to us and pointing out further interesting connections between group theory and homotopy theory.

1 Proof of corollaries

It is known that the generalized continuum hypothesis ensures the existence of arbitrarily large cardinals λ which satisfy Dl_λ . For instance, successor cardinals $\lambda = \kappa^+$ of regular cardinals $\kappa > \aleph_0$ satisfy Dl_λ . Hence, in order to prove Corollary 3, we next show that the inclusion $\eta : H \hookrightarrow \mathcal{G}$ given in the Main Theorem is a localization. Suppose that $\varphi : H \rightarrow \mathcal{G}$ is a homomorphism. We have to show that there is a unique homomorphism

$\Phi : \mathcal{G} \rightarrow \mathcal{G}$ such that $\Phi\eta = \varphi$. If $\varphi = 0$ then $\Phi = 0$ makes the diagram (0.1) commutative. To see that it is unique, we note that H is in the kernel K of Φ and $K = \mathcal{G}$ by simplicity, hence $\Phi = 0$. Now suppose that $\varphi \neq 0$. Since H is simple we have that $\varphi(H) \leq \mathcal{G}$ and $H \cong \varphi(H)$. Thus by part (b) of the Main Theorem there is an element $y \in \mathcal{G}$ such that $\varphi = y^* \upharpoonright H$ where y^* is the inner automorphism of $\mathcal{G}$ given by $y^*(x) = y^{-1}xy$ for all $x \in \mathcal{G}$, and $y^* \upharpoonright H$ denotes its restriction on H . Hence $\Phi = y^*$ satisfies $\Phi\eta = \varphi$. Suppose that $\Phi' : \mathcal{G} \rightarrow \mathcal{G}$ is another homomorphism such that $\Phi'\eta = \varphi$. Then $\Phi' \neq 0$. Now since $\mathcal{G}$ is simple by (f) and co-hopfian by (e) we have that Φ' is an automorphism. Thus, by completeness (d) we have $\Phi' = g^*$ for some $g \in \mathcal{G}$. Now $(g^{-1}y)^* : \mathcal{G} \rightarrow \mathcal{G}$ fixes all elements of H . Hence, by part (c), we finally obtain that $g^{-1}y = 1$ and thus $\Phi = \Phi'$ as desired.

Recall from [5] or [3] that a map $f : X \rightarrow Y$ between two connected spaces is a *homotopical localization* if Y is f -local, i.e. if the map of pointed function spaces

$$\mathrm{map}_*(Y, Y) \rightarrow \mathrm{map}_*(X, Y)$$

induced by composition by f is a weak homotopy equivalence. As in the case of groups this says that $Y \simeq L_f X$, where L_f is the localization functor with respect to f .

It turns out that the homotopical localizations of the circle $S^1 = K(\mathbb{Z}, 1)$ are precisely Eilenberg–Mac Lane spaces $K(A, 1)$ where A ranges over the class of all E -rings [3, Theorem 5.11]. And therefore this class is proper (i.e. it is not a set) in view of the result in [7]. Recall that an E -ring is a commutative ring A with identity which is canonically isomorphic to its own ring of additive endomorphisms. Corollary 4 claims that a similar situation holds for $K(H, 1)$ if H is a suitable group. However, in this case, other localizations of $K(H, 1)$ which are not of the form $K(G, 1)$ may exist.

Corollary 4 follows from the fact that every localization of groups $H \rightarrow \mathcal{G}$ gives rise to a localization of spaces $K(H, 1) \rightarrow K(\mathcal{G}, 1)$. This holds because, for arbitrary groups A and B , the space $\mathrm{map}_*(K(A, 1), K(B, 1))$ is homotopically discrete and equivalent to the set $\mathrm{Hom}(A, B)$.

We recommend the reader not so familiar with the basic facts on homotopy theory to consult, for instance, [18] or the monograph by Aubry [1] (lectures of a DMV seminar by Baues, Halperin and Lemaire) as well as Dror Farjoun’s exposition [5] for homotopical localizations.

2 Sketch of the proof of Main Theorem

The group $\mathcal{G}$ in our main theorem has cardinality λ^+ , where λ^+ is the successor cardinal of a regular cardinal λ . It will be the union of a λ^+ -filtration $\{\mathcal{G}_\alpha : \alpha < \lambda^+\}$, that is,

$$\mathcal{G} = \bigcup_{\alpha < \lambda^+} \mathcal{G}_\alpha$$

where $\mathcal{G}_0 = \text{Aut}(H)$ and $|\mathcal{G}_\alpha| = \lambda$ for $0 \neq \alpha < \lambda^+$. Moreover $|\mathcal{G}_{\alpha+1} \setminus \mathcal{G}_\alpha| = \lambda$ as well. More precisely, at each step α , $\mathcal{G}_\alpha$ will be the union of a certain directed system $\mathbb{G}_\alpha$ of size $|\mathbb{G}_\alpha| = \lambda$, where each subgroup $G \in \mathbb{G}_\alpha$ has cardinality $< \lambda$. It is convenient to give a linear order to all elements which are added in the construction of $\mathcal{G}$, so we view any $G \in \mathbb{G}_\alpha$ as a subset of

$$\widetilde{\lambda^+} := \{\lambda\alpha' + i : \alpha' \in \lambda^+, i < \lambda\}.$$

This set equals $\lambda^+ \times \lambda$ with the lexicographical order on the pair of ordinals, hence it is well-ordered and has cardinality $|\widetilde{\lambda^+}| = \lambda^+$. Hence $\widetilde{\lambda^+}$ will be the universe for $\mathcal{G}$. All groups $G \in \mathbb{G}_\alpha$ ($0 \neq \alpha < \lambda^+$) –which can be thought as “approximations” to $\mathcal{G}$ – will satisfy properties (a), (b) and (c) of the Main Theorem, and moreover $G \subseteq \lambda\zeta_\alpha$ for some unbounded, strictly increasing sequence of ordinals $\zeta_\alpha < \lambda^+$. The construction of these approximations is given in Section 4 (only as groups). We show in Section 6 how they can be embedded into $\widetilde{\lambda^+}$; see Proposition 6.4. The group structure of each G is coded, up to a certain strong isomorphism, by a pair $p = (\gamma, u)$ with $\gamma \in \lambda$ and $u \subset \lambda^+$ of cardinality $|u| < \lambda$ (u is the image of G under the natural projection $\widetilde{\lambda^+} \rightarrow \lambda^+$). All such pairs p are collected in a partially ordered set $\mathbb{P}$ which satisfies the requirements of the $\mathbb{P}$ -game defined by Hart, Laflamme and Shelah in [9], which we have included in Appendix A for convenience. A “local” construction taking care of simplicity (f) is done in Proposition 4.14. The properties to be complete (d) and co-hopfian (e) are treated together in Section 5 and again in Section 7 inside the universe λ^+ . Finally, in Section 8 we explain how the two players of the $\mathbb{P}$ -game carry out the “global” construction of $\mathcal{G}$: player I defines adequate “density systems” which make some restrictions to player II in his choice of the sequence ζ_α and the directed systems $\mathbb{G}_\alpha$. These restrictions will guarantee the desired properties of $\mathcal{G}$. The combinatorial principle Dl_λ , which is a weak version of diamond $\diamond(\lambda)$, is used by player II to “predict” in some sense the moves of player I.

3 Some preliminaries on set theory, free products with amalgamation and HNN extensions

We recall some basic notions of set theory from [10].

The cofinality $\text{cf}(\lambda)$ of an infinite cardinal λ is the minimal cardinal κ for which there exists an increasing sequence of ordinals $\lambda_i < \lambda$ of length κ such that $\sup\{\lambda_i : i < \kappa\} = \lambda$. Clearly, $\text{cf}(\lambda) \leq \lambda$. A cardinal λ is called *regular* if $\text{cf}(\lambda) = \lambda$.

Let λ be an uncountable regular cardinal. A subset $C \subseteq \lambda$ is a *club* if it is closed under limits of length $< \lambda$ and if it is unbounded. A subset $S \subseteq \lambda$ is *stationary* if $S \cap C \neq \emptyset$ for all clubs C in λ . In particular every club is stationary. Let λ^+ be the successor cardinal of λ . We will use that $C = \{\delta < \lambda^+ : \text{cf}(\delta) = \lambda\}$ is a club of λ^+ . Moreover, to prove that the group $\mathcal{G}$ is complete (Section 9.4) we need Fodor's Lemma, see Jech [10, p. 59, Theorem 22].

Lemma 3.1 *Suppose $\eta : S' \rightarrow \lambda^+$ is a regressive function where S' is a stationary subset of λ^+ . Then there is a stationary $S \subseteq S'$ such that η is constant on S .*

Recall that the generalized continuum hypothesis GCH says that for any cardinal κ the successor κ^+ coincides with $|\mathcal{P}(\kappa)| = 2^\kappa$, where $\mathcal{P}(\kappa)$ denotes the set of subsets of κ . Moreover GCH implies that $|\mathcal{P}_{<\lambda}(\lambda)| = \lambda^{<\lambda}$ is λ , where $\mathcal{P}_{<\lambda}(\lambda)$ denotes the set of subsets of λ of cardinality $< \lambda$.

In the next section we shall need two lemmas describing torsion subgroups of free products with amalgamations and HNN extensions (compare with Theorem 2.7, p. 187, Theorem 6.6, p. 212 and Theorem 2.4, p. 185 in [13]).

Lemma 3.2 *Let $G^* = G *_{{G_0}} G'$ be the free product of G and G' amalgamating a common subgroup G_0 . If H' is a torsion subgroup of G^* then there exists a $y \in G^*$ such that $(H')^y$ is contained in G or in G' .*

PROOF. Suppose H' is not contained in G_0 , otherwise we are done. Fix $h \in H' \setminus G_0$ and let h be written in a canonical form

$$h = g_1 g'_1 \cdots g_n g'_n,$$

i.e. $g_i \in G \setminus G_0$ and $g'_i \in G' \setminus G_0$ (with the convention that g_1 or g'_n may be trivial.) Note that this expression is not unique in general, but its length $|h|$ only depends on h .

Suppose first that all elements of H' have length 1. If H' contains two elements $g \in G \setminus G_0$ and $g' \in G' \setminus G_0$ then $gg' \in H$ is a word written in a reduced normal form, and has length 2, hence it is non torsion by Theorem 2.6, p. 187 in [13]. This contradicts that H is torsion. Thus, all $g \in H'$ must belong to the same factor, and we are done.

We can suppose that the first factor g_1 of h is non trivial and that $|h|$ is odd and ≥ 3 . Choose h such that $|h|$ is the minimum over all lengths of elements in H' of length ≥ 3 . If $g'_n \neq 1$ then we obtain a contradiction as before. Thus g'_n must be trivial and similarly $g_n g_1 \in G_0$. Therefore, h has a normal decomposition of the form

$$h = g_1 g'_1 \cdots \bar{g}'_{n-1} g_1^{-1},$$

for some $\bar{g}'_{n-1} \in G' \setminus G_0$ which we rename by g'_{n-1} to simplify the notation.

Similarly, all $h_1 \in H'$ must have odd length. Moreover, if $|h_1| \geq 3$ then h_1 has a normal form starting with g_1 , the first factor of h and ending with g_1^{-1} . Of course, if $|h_1|=1$ then $h_1 \in G$.

Consider then $g_1^{-1} H' g_1$. If $h_1 \in H'$ has length ≥ 3 , then $g_1^{-1} h_1 g_1$ has length $|h| - 2$, and its normal form begins and ends with elements in G' . On the other hand, if $h_1 \in H'$ has length 1, then $g_1^{-1} h_1 g_1 \in G$. But, arguing as before, we see that such elements of length 1 must be actually in G_0 .

Repeating this process we obtain a $y \in G^*$ such that $h = ygy^{-1}$ where $g \in G \setminus G_0$ or $G' \setminus G_0$ depending on n , if n is odd or even, respectively. If any of the elements in $(H')^y$ has length ≥ 3 , then we can conclude that $g \in G_0$ which is a contradiction. Thus all elements in $(H')^y$ must have length 1, and the first case applies, hence $(H')^y$ is contained in some factor. This finishes the proof. $\square$

Lemma 3.3 *Let G be any group, and $\phi : G_0 \rightarrow G_1$ be an isomorphism between two subgroups of G . Consider the HNN extension $G^* = \langle G, t : t^{-1}ht = \phi(h), h \in G_0 \rangle$. If H' is a torsion subgroup of G^* , then there exists a $y \in G^*$ such that $(H')^y$ is contained in G .*

PROOF. The proof is similar to that of Lemma 3.2. $\square$

The following lemma describes centralizers of torsion subgroups in free products with amalgamation. An analogous lemma for HNN extensions also holds.

Lemma 3.4 *Let $G^* = G *_{G_0} G'$ be the free product of G and G' amalgamating a common subgroup G_0 . Let $H' \leq G$ be a torsion subgroup and let $1 \neq x \in G^*$ an element which commutes with all elements of H' . Then either $x \in G$ or $(H')^g \leq G_0$ for some $g \in G^*$.*

PROOF. Let $H' \leq G$ be a torsion subgroup, and $x \in G^*$ such that $[x, H'] = 1$. If $x \in G$ we are done. Suppose then that $x \notin G$. Express x in a reduced normal form

$$x = g_1 g'_1 \cdots g_n g'_n,$$

that is, $g_i \in G \setminus G_0$ and $g'_i \in G' \setminus G_0$, where $g'_1 \neq 1$ (g_1 or g'_n may be trivial). The relation $h^{-1} x^{-1} h x = 1$ yields the following

$$h^{-1} g'^{-1}_n g_n^{-1} \cdots g'^{-1}_1 (g_1^{-1} h g_1) g'_1 \cdots g_n g'_n = 1.$$

By the normal form theorem for free products with amalgamation [13, Theorem 2.6 p 187], this is only possible if $g_1^{-1} h g_1 \in G_0$ for all $h \in H'$. This concludes the proof.

□

4 Approximations to $\mathcal{G}$

In this section we fix a suitable group H , we denote $\widehat{H} = \text{Aut}(H)$ for shortness and view $H \leq \widehat{H}$ as subgroup. We also fix an uncountable regular cardinal λ such that $|\widehat{H}| < \lambda$. As usual $C_{G'}(G)$ and $N_{G'}(G)$ denote, respectively, the centralizer and the normalizer of a subgroup G in a group G' .

We next consider two particular families $\mathcal{K}^* \subseteq \mathcal{K}$ of groups which will be used to approximate our group $\mathcal{G}$ of the Main Theorem in the sense explained in Section 2.

First we prove a series of lemmas which culminate in Proposition 4.11. This states that the class of groups $\mathcal{K}^*$ is *dense* in $\mathcal{K}$, that is, given an infinite group $G \in \mathcal{K}$ there is a group $G' \in \mathcal{K}^*$ of the same cardinality such that $G \leq G'$. In particular, we provide an explicit algorithm to construct groups satisfying conditions (a), (b) and (c) of the Main Theorem.

We also show Proposition 4.14 which will be used to prove that our final group $\mathcal{G}$ is simple.

Definition 4.1 1. $\mathcal{K}$ consists of all groups G with $|G| < \lambda$ such that $H \leq \widehat{H} \leq G$, and any isomorphic copy of H in G has trivial centralizer in G . That is,

$$\mathcal{K} = \{G : \widehat{H} \leq G, |G| < \lambda, \text{ if } H \cong H' \leq G, x \in G \text{ with } [H', x] = 1, \text{ then } x = 1\},$$

where $[x, y] = x^{-1}y^{-1}xy$.

2. $\mathcal{K}^*$ is the class of all groups G in $\mathcal{K}$ which are generated by copies of H and such that any isomorphism between H and a subgroup of G is induced by conjugation with an element in G .

Note that condition (1) in Definition 4.1 is stronger than property (c) of the Main Theorem. If we denote by $H(G)$ the H -socle of G , i.e. the subgroup of G generated by all images of homomorphisms from H into G , then $H(G) = G$ for all $G \in \mathcal{K}^*$. We start observing that $\mathcal{K}$ is not empty. This is obvious since H is suitable and $|\widehat{H}| < \lambda$.

Lemma 4.2 $\widehat{H}$ is in $\mathcal{K}$. $\square$

Lemma 4.3 If G and G' are in $\mathcal{K}$ then $G * G' \in \mathcal{K}$.

PROOF. Suppose that $H' \leq G * G'$ with $H' \cong H$ and $x \in G * G'$ such that $[H', x] = 1$. By Lemma 3.2, we can suppose that $(H')^y \leq G$. Hence $[H', x] = 1$ implies $[(H')^y, x^y] = 1$. In other words,

$$(h^y)^{-1}(x^y)^{-1}h^yx^y = 1 \tag{4.1}$$

for all $h \in H'$. If $x^y \notin G$, then express $(x^y)^{-1}$ in reduced form and replace a first choice $h \neq 1$ by a different one if the first G -factor of $(x^y)^{-1}$ in (4.1) is cancelled by $(h^y)^{-1}$. Hence the normal form of the left hand in (4.1) has non-trivial factors and the normal form theorem for free products [13, p. 175] gives a contradiction, hence $x^y \in G$. But $G \in \mathcal{K}$, hence $x^y = 1$ and thus $x = 1$, as desired. Note that the lemma also follows from Lemma 3.4. $\square$

Lemma 4.6 below will take care of part (b) of the Main Theorem. If G is any group in $\mathcal{K}$ and $\phi : A \rightarrow B$ is an isomorphism between two subgroups of G isomorphic to H , we want that ϕ is the restriction of an inner automorphism in some extension $G \leq G_2 \in \mathcal{K}$. This can be established using HNN extensions as we next show. We will get this after several steps.

Lemma 4.4 *Let $G \in \mathcal{K}$ and $B \leq G$ be a subgroup isomorphic to H . Then there is $G \leq G_1 \in \mathcal{K}$ such that $\text{Aut}(B) \leq G_1$.*

PROOF. Let $\widehat{B} = \text{Aut}(B)$ and $N = N_G(B)$. If $\widehat{B} \leq G$ then $G_1 = G$. Suppose that $\widehat{B} \not\leq G$. Note that $N = G \cap \widehat{B}$ so we can consider the free product with amalgamation $G_1 := G *_N \widehat{B}$. We shall show that $G_1 \in \mathcal{K}$. Let $H' \leq G_1$ be a subgroup isomorphic to H and $1 \neq x \in G_1$ such that $[H', x] = 1$. By Lemma 3.2 we can suppose that $H' \leq G$ or $H' \leq \widehat{B}$. Suppose that $H' \leq G$, the other case is easier. Let $x = g_1 g_2 \cdots g_n$ be written in a reduced normal form. If $x = g_1 \in G$ then $x = 1$ since $G \in \mathcal{K}$, and this is a contradiction. Hence $x = g_1 \in \widehat{B} \setminus N$. As in Lemma 3.4 we deduce that $H' = (H')^{g_1} \leq N$, thus $H' = B$ since B is suitable (condition 2). Hence $g_1 \in N$ and we get a contradiction. If $n = 2$, then we obtain $(H')^{g_1} = (H')^{g_2} = B = H'$. So both g_1 and g_2 are in N , which is a contradiction. Similarly, if $n \geq 3$ we have that g_2 and g_3 are in N . This is again impossible. This concludes the proof. $\square$

By the previous lemma we can suppose that if $B \leq G \in \mathcal{K}$, and if $B \cong H$, then $\widehat{B} \leq G$ as well.

Recall that two subgroups A and B of a group G are conjugate in G if there is $g \in G$ such that $A = B^g$. In particular, if $A, B \leq G$, $A \cong B \cong H$ and $\widehat{A}, \widehat{B}$ are conjugate in G then A and B are also conjugate. Indeed, if $g \in G$ such that $g^* : \widehat{A} \cong \widehat{B}$, then the image $A^g \leq \widehat{B}$ is a subgroup isomorphic to B , hence $A^g = B$, by condition 2 of Definition 1.

Lemma 4.5 *Let $G \in \mathcal{K}$ and $B \leq \widehat{B} \leq G$, where $B \cong H$. Suppose that H and B are not conjugate in G . Let $\phi : \widehat{H} \rightarrow \widehat{B}$ be any isomorphism. Then, the HNN extension*

$$G_1 = \langle G, t : t^{-1}ht = \phi(h) \text{ for all } h \in \widehat{H} \rangle$$

is also in $\mathcal{K}$.

PROOF. Let $H' \leq G_1$ be a subgroup isomorphic to H' and $1 \neq x \in G_1$ such that $[H', x] = 1$. By Lemma 3.3 we can suppose that $H' \leq G$ already. Let

$$x = g_0 t^{\varepsilon_1} g_1 t^{\varepsilon_2} \cdots g_{n-1} t^{\varepsilon_n} g_n$$

be written in a reduced form in G_1 , where $g_i \in G$ and there is no subwords $t^{-1}g_i t$ with $g_i \in \widehat{H}$ or $tg_i t^{-1}$ with $g_i \in \widehat{B}$ (see [13, p. 181]).

If $x = g_0 \in G$ then $x = 1$, since $G \in \mathcal{K}$. This yields a contradiction. Thus $n \geq 1$. We have $[h, x] = h^{-1}x^{-1}hx = 1$ for every $h \in H'$. In other words, for a fixed $h \neq 1$, the following holds

$$h^{-1}g_n^{-1}t^{-\varepsilon_n} \dots t^{-\varepsilon_1}(g_0^{-1}hg_0)t^{\varepsilon_1} \dots t^{\varepsilon_n}g_n = 1. \quad (4.2)$$

By the normal form theorem for HNN extensions ([13, p. 182]), either $\varepsilon_1 = 1$ and $g_0^{-1}hg_0 \in \widehat{H}$, or $\varepsilon_1 = -1$ and $g_0^{-1}hg_0 \in \widehat{B}$. Suppose that $\varepsilon_1 = 1$, the other case is analogous. Then $(H')^{g_0} \leq \widehat{H}$, thus $(H')^{g_0} = H$, and we can replace in (4.2) the subword $t^{-1}(g_0^{-1}hg_0)t$ by $\phi(g_0^{-1}hg_0) \in B$. Repeating the same argument we obtain that $\varepsilon_i = 1$ or -1 . Hence one of the two possibilities holds depending on $\varepsilon_2 = 1$ or $\varepsilon_2 = -1$. So we have either

$$\text{id} : H_1 \xrightarrow{g_0^*} \widehat{H} \xrightarrow{\phi} \widehat{B} \xrightarrow{g_1^*} \widehat{H} \xrightarrow{\phi} \dots \xrightarrow{g_n^*} H_1$$

or

$$\text{id} : H_1 \xrightarrow{g_0^*} \widehat{H} \xrightarrow{\phi} \widehat{B} \xrightarrow{g_1^*} \widehat{B} \xrightarrow{\phi^{-1}} \dots \xrightarrow{g_n^*} H_1.$$

In the first case we have an isomorphism $g_1^*\phi : \widehat{H} \xrightarrow{\phi} \widehat{B} \xrightarrow{g_1^*} \widehat{H}$. Since $\widehat{H}$ is complete there is $g \in \widehat{H}$ such that $g_1^*\phi = g^*$. This yields $\phi = (g_1^{-1}g)^*$, i.e. $\widehat{H}$ and $\widehat{B}$ are conjugate, and thus H and B are conjugate, but this is impossible by hypothesis. In the second case we have $g_1 \in \widehat{B}$ by completeness. But, on the other hand $g_1 \notin \widehat{B}$ since x is written in a reduced form. We conclude that G_1 is in $\mathcal{K}$, as desired. $\square$

Let $\text{pInn}(G)$ denote the set of *partial inner automorphisms*, which are the isomorphisms $\phi : G_1 \rightarrow G_2$ where $G_1, G_2 \leq G$ such that ϕ can be extended to an inner automorphism of G . Hence $\text{pInn}(G)$ are all restrictions of conjugations to subgroups of G .

Lemma 4.6 *Let $G \in \mathcal{K}$. Let $A \leq \widehat{A} \leq G$ and $B \leq \widehat{B} \leq G$. Let $\phi : A \cong B$ be any isomorphism. Then, there is $G_2 \in \mathcal{K}$ such that $\phi \in \text{pInn}(G_2)$. Moreover, G_2 can be obtained from G by at most two successive HNN extensions.*

PROOF. If $\phi \in \text{pInn}(G)$ we take $G_2 = G$. Suppose that $\phi \notin \text{pInn}(G)$. If $\widehat{H}$ and $\widehat{A}$ are conjugate, we take $G_1 = G$. Otherwise, we consider the HNN extension

$$G_1 = \langle G, t_1 : t_1^{-1}ht_1 = \phi_1(h) \text{ for all } h \in \widehat{H} \rangle$$

where ϕ_1 is any isomorphism between $\widehat{H}$ and $\widehat{A}$. By Lemma 4.5 we know that $G_1 \in \mathcal{K}$. Now, if H and B are conjugate in G_1 , we take $G_2 = G_1$. It follows automatically that

$\phi \in \text{pInn}(G_1)$, since $\widehat{H}$ is complete. If $\widehat{H}$ and $\widehat{B}$ are not conjugate in G_1 , we consider a new HNN extension

$$G_2 = \langle G_1, t_2 : t_2^{-1} h t_2 = \phi_2(h) \text{ for all } h \in \widehat{H} \rangle$$

where ϕ_2 is any isomorphism between $\widehat{H}$ and $\widehat{B}$. Again $G_2 \in \mathcal{K}$ by Lemma 4.5. In that case we have an isomorphism $(t_2^{-1})^* \phi t_1^* : \widehat{H} \rightarrow \widehat{H}$, which equals g^* for some $g \in \widehat{H}$ by completeness. Thus $\phi = (t_2 g t_1^{-1})^* \upharpoonright A$. This shows that $\phi \in \text{pInn}(G_2)$, as desired. $\square$

The following two lemmas will be used to prove Lemma 4.10. This will take care of part (a) of the Main Theorem. Let $o(g)$ denote the order of $g \in G$.

Lemma 4.7 *Let $G \in \mathcal{K}$ and suppose that $G' \in \mathcal{K}$ or G' does not contain any subgroup isomorphic to H . Let $g \in G$ and $g' \in G'$ with $o(g) = o(g')$. Then $(G * G')/N \in \mathcal{K}$ where N is the normal subgroup of $G * G'$ generated by $g^{-1}g' \in G * G'$.*

PROOF. Let $\overline{G} = (G * G')/N$ and view G and G' as subgroups of $\overline{G}$, respectively. Suppose that we have a subgroup $H' \leq \overline{G}$ isomorphic to H and $x \in \overline{G}$ such that $[H', x] = 1$. By Lemma 3.2 a conjugate of H' is contained in G or G' . Hence by hypothesis we can assume that H' is already contained in G . Suppose that $x \neq 1$. By Lemma 3.4 it follows that either $x \in G$ or a conjugate of H' is contained in $\langle g \rangle$. In the first case $x = 1$ since $G \in \mathcal{K}$, so contradiction, and the second case is obviously impossible. Thus $\overline{G} \in \mathcal{K}$ as desired. $\square$

Lemma 4.8 *If $g \in G \in \mathcal{K}$, then there is a group $\overline{G} \in \mathcal{K}$, such that $G \leq \overline{G}$, with $|\overline{G}| = |G| \cdot \aleph_0$ and $g \in H(\overline{G})$.*

PROOF. Suppose that $o(g) = \infty$ and that $g \notin H(G)$. Let H_1 and H_2 be two isomorphic copies of H . Choose a non trivial element $h \in H$ and let h_1 and h_2 be its copies in H_1 and H_2 respectively. Now define

$$\overline{G} = (G * H_1 * H_2)/N$$

where N is the normal subgroup generated by $g^{-1}h_1h_2$. Then $\overline{G} \in \mathcal{K}$ by Lemma 4.7 and moreover $g \in H(\overline{G})$.

If $o(g) = n < \infty$ we first embed $G \leq (G * K)/N$ where $K = \langle x_1, x_2 : (x_1 x_2)^n = 1 \rangle$ and N is the normal closure of $g^{-1} x_1 x_2$. Then, by the previous lemma $(G * K)/N \in \mathcal{K}$. Now, since $o(x_1) = o(x_2) = \infty$, we can apply the first case. $\square$

We next show that the class $\mathcal{K}$ is closed under limits of relatively “small” length.

Lemma 4.9 *Let $\gamma < \lambda$ and $\{G_i : i < \gamma\}$ be an ascending continuous chain of groups in $\mathcal{K}$. Then the union $G = \bigcup_{i < \gamma} G_i$ also belongs to $\mathcal{K}$.*

PROOF. Let $H' \leq G$ be a subgroup isomorphic to H and let $x \in G$ be such that $[H', x] = 1$. Since $|H'| < \lambda$ and λ is regular we have $H' \leq G_{i_0}$ for some index i_0 . We can also suppose that $x \in G_{i_0}$. Now $G_{i_0} \in \mathcal{K}$, hence $x = 1$. $\square$

Lemma 4.10 *If $G \in \mathcal{K}$ then there is a group $G' \in \mathcal{K}$ such that $G \leq G'$ with $|G'| = |G| \cdot \aleph_0$ and $H(G') = G'$.*

PROOF. By Lemma 4.3 we may assume that G is infinite, hence the last cardinal condition becomes simply $|G'| = |G|$. Applying Lemma 4.8 $|G|$ -times and Lemma 4.9 once we can find an equipotent group $G_1 \in \mathcal{K}$, such that $G \leq G_1$ and $G \leq H(G_1)$, inductively we find $G \leq G_{n+1} \in \mathcal{K}$ such that $G_n \leq H(G_{n+1})$ for each natural $n \geq 1$. The union G' of this chain belongs to $\mathcal{K}$ by Lemma 4.9 and obviously satisfies $H(G') = G'$. $\square$

For example, using Lemma 4.3 and Lemma 4.9 we obtain that, for every cardinal $\kappa < \lambda$, the free product $*_{\alpha \in \kappa} \widehat{H}_\alpha$ belongs to $\mathcal{K}$, where $\widehat{H}_\alpha$ is an isomorphic copy of $\widehat{H}$ for every $\alpha \in \kappa$. Now we can apply the following proposition to obtain a group $G' \in \mathcal{K}^*$, such that $*_{\alpha \in \kappa} \widehat{H}_\alpha \leq G'$.

Proposition 4.11 *$\mathcal{K}^*$ is dense in $\mathcal{K}$.*

PROOF. Let $G \in \mathcal{K}$. We have to construct a group G' in $\mathcal{K}^*$ such that $G \leq G'$ with $|G'| = |G| \cdot \aleph_0$. This can be obtained easily by using alternatively Lemma 4.6 and Lemma 4.10 a countable number of times. The limit group will still lie in $\mathcal{K}$ by Lemma 4.9. Moreover it is in $\mathcal{K}^*$ by construction. $\square$

We remark that $\mathcal{K}^*$ is also closed under direct limits of relatively small length. The proof is similar to that of Lemma 4.9.

Lemma 4.12 *Let $\gamma < \lambda$ and let $\{G_i : i < \gamma\}$ be an ascending continuous chain of groups in $\mathcal{K}^*$. Then the union $G = \bigcup_{i < \gamma} G_i$ also belongs to $\mathcal{K}^*$. $\square$*

Lemma 4.13 *Let $G^* = G *_{G_0} G'$ be the free product of G and G' amalgamating a common subgroup G_0 . Suppose that G , G' and G_0 are in $\mathcal{K}^*$. Then $G^* \in \mathcal{K}$.*

PROOF. Let $H' \leq G^*$ be a subgroup isomorphic to H , and $x \in G^*$ such that $[H', x] = 1$. As in previous results we can assume that $H' \leq G_0$ and that $x = g_1 g_2 \cdots g_n$, written in a reduced form, has length bigger than two. Then we have $g_1^* : H' \cong (H')^{g_1}$ both of them inside G_0 . Since $G_0 \in \mathcal{K}^*$ there exists a $g \in G_0$ such that $g^* : H' \cong (H')^{g_1}$. We can also suppose that the automorphism group $\widehat{H}'$ is already in G_0 , by Lemma 4.4. Hence, the composition $(g_1^{-1}g)^* : H' \rightarrow H'$ is an automorphism, which is inner by completeness. Thus, $g_1^{-1}g \in G_0$ and thus $g_1 \in G_0$. This is a contradiction, since x was written in a reduced form. $\square$

In order to show that our final group $\mathcal{G}$ is simple we only must consider normal subgroups N of $\mathcal{G}$ which are cyclically generated, i.e. there is an $1 \neq x \in \mathcal{G}$ with $N = \langle x^{\mathcal{G}} \rangle$. We need that $N = \mathcal{G}$. There are two natural cases depending on the order of x . The case that x has infinite order is taken care by the next Proposition 4.14. Hence assuming that all elements of infinite order are conjugate, a consequence of Proposition 4.14, we only need to note that any element of finite order can be written as a product of two elements of infinite order, just take y from a different factor then $g = (gy)y^{-1}$. Hence $\mathcal{G} = N$. If x has finite order, then there is a conjugate y of x such that xy has infinite order. Hence $xy \in N$ and the first case applies. In Proposition 8.3 we shall also use this argument.

Proposition 4.14 *Let G be a group in $\mathcal{K}$. Let $g, f \in G$, where $o(f) = o(g) = \infty$ and g does not belong to the normal subgroup generated by f . Then there is a group $\overline{G} \in \mathcal{K}$ such that $G \leq \overline{G}$ and g is conjugate to f in $\overline{G}$.*

PROOF. Let $\alpha : \langle f \rangle \rightarrow \langle g \rangle$ be the isomorphism mapping f to g . By hypothesis $\alpha \notin \text{pInn } G$. As in Lemma 4.6 consider the HNN extension $\overline{G} = \langle G, t : t^{-1}ft = g \rangle$. We must show that $\overline{G} \in \mathcal{K}$. Clearly $|\overline{G}| < \lambda$ and consider any H' with $H \cong H' \leq G$ and any $x \in \overline{G}$ with $[H', x] = 1$. By the argument in Lemma 4.5 we may assume that $H' \leq G$ and $x \in \overline{G}$ with $[H', x] = 1$. Now we apply Lemma 3.4 as we did in Lemma 4.5. $\square$

5 Small cancellation groups

Let H be as in Section 4. Let $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$ be the group of outer automorphisms of a group G . We say that G is *complete* if $\text{Out}(G) = 1$ and if the centre $Z(G) = 1$, see Robinson [14] for instance. Furthermore, G is called *co-hopfian* if every monomorphism from G to G is an automorphism.

In this section we shall use small cancellation theory as in Shelah [16] or Schupp [15] with the purpose of “killing” monomorphisms which are not inner automorphisms. This will yield that $\mathcal{G}$ is complete and co-hopfian. Recall from [15] that an automorphism ψ of a group G_0 is inner if and only if whenever G_0 is embedded into a group G then ψ extends to some automorphism of G .

An alternative tool would be the use of wreath products, as in [6] where Dugas and Göbel answered a question of Hall about the existence of particular complete groups.

Let us first draw a picture which will appear infinitely many times along the construction of $\mathcal{G}$:

$$\begin{array}{ccccc}
 & & G^* & & \\
 & \nearrow & \downarrow & \nwarrow & \\
 x_0 \in H_0 \subset G_1 & \rightarrow & G & \leftarrow & G_2 \supset H_1 \ni x_1 \\
 & \nwarrow & & \nearrow & \\
 & & z \in G_0 & &
 \end{array} \tag{5.1}$$

Here, $G_0 \leq G_1$, $G_0 \leq G_2$ are all groups in $\mathcal{K}$ and $G^* = G_1 *_{G_0} G_2$. For $i = 0, 1$ we have subgroups $H_i \leq G_{i+1}$ isomorphic to H such that $H_i \cap G_0 = \{1\}$, and elements $1 \neq x_i \in H_i$, $z \in G_0$.

Our task now is to find a new group G in $\mathcal{K}$ where $G_0 \leq G$ such that:

- (1) $z = \tau(x_0, x_1)$ in G for some word $\tau(x_0, x_1)$ depending on the letters x_0 and x_1 .
- (2) G contains G_1 and G_2 as malnormal subgroups, i.e. for every element $y \in G \setminus G_i$ we have $(G_i)^y \cap G_i = \{1\}$.
- (3) Suppose that $\psi_0 : G_0 \rightarrow G_0$ is a monomorphism which extends to a monomorphism ψ with domain G . Suppose that $\psi \upharpoonright H_i = y_i^* \upharpoonright H_i$ with $y_i \in G_{i+1} \setminus C_{G_{i+1}}(x_i)$ for $i = 0, 1$. Then $y_0, y_1 \in G_0$.

If ψ is a monomorphism of our final group $\mathcal{G}$ which is not an inner automorphism then we will deal with this map restricted to many subgroups G of $\mathcal{G}$ which are in $\mathcal{K}^*$. On these

restrictions $\psi \upharpoonright G$ we will recognize that $\psi \upharpoonright G$ is not inner, by the following result. Call a monomorphism of a group λ -inner if it acts by conjugation on all subgroups generated by $< \lambda$ elements. In particular $\aleph_0$ -inner is locally inner in the classical sense.

Proposition 5.1 *Suppose that our final group $\mathcal{G}$ is constructed as a direct limit of groups in $\mathcal{K}^*$ and $\mathcal{G} = H(\mathcal{G})$. If ψ is a monomorphism of $\mathcal{G}$ which is λ -inner, then ψ is an inner automorphism.*

PROOF. Since $\mathcal{G}$ satisfies all conditions for groups in $\mathcal{K}^*$ except that $|\mathcal{G}| = \lambda^+$, we know that $H \leq \mathcal{G}$ and $\psi(H)$ lies in some subgroup which is in $\mathcal{K}^*$ hence $\psi \upharpoonright H$ is conjugation by some element in $\mathcal{G}$. Hence we can suppose without loss of generality that $\psi \upharpoonright H = \text{id}$.

Now let $H' \leq \mathcal{G}$ be another copy of H . As before there is an element $y \in \mathcal{G}$ such that $\psi \upharpoonright H' = y^* \upharpoonright H'$. The subgroup H' and y generate a subgroup $G \leq \mathcal{G}$ which is generated by $< \lambda$ elements, and it follows from properties of $\mathcal{G}$ that we may also assume $G \in \mathcal{K}^*$. By hypothesis on ψ there is an element $g \in \mathcal{G}$ such that $\psi \upharpoonright HH' = g^* \upharpoonright HH'$. Therefore, for all $h \in H$ and $h' \in H'$, we will have

$$hy^{-1}h'y = \psi(h)\psi(h') = \psi(hh') = g^{-1}hh'g.$$

In particular, if $h' = 1$, then $g^{-1}hg = h$ for all $h \in H$. Since $G \in \mathcal{K}$ it follows that $g = 1$. Thus $h' = y^{-1}h'y$ for all $h' \in H'$. By the same reason $y = 1$. We have shown that $\psi \upharpoonright H' = \text{id}$ for all $H' < \mathcal{G}$. Since $H(\mathcal{G}) = \mathcal{G}$, we conclude $\psi = \text{id}$. $\square$

Here we give an argument which will provide extensions G of G_0 which do not allow extensions of $\psi \upharpoonright G_0$. However G will be build into $\mathcal{G}$, hence the total map ψ is a contradiction.

More technically, the relations given by (1) will ensure that z can only be moved by conjugation. This will prove that $\mathcal{G}$ is complete and co-hopfian at the same time.

The following “long” and “complicated” word in x_0 and x_1 is a good candidate for our purposes:

$$\tau(x_0, x_1) = (x_0x_1)(x_0x_1^2)(x_0x_1)^2(x_0x_1^2)^2 \cdots (x_0x_1)^{80}(x_0x_1^2)^{80}. \quad (5.2)$$

Then define

$$r(z, x_0, x_1) := z^{-1}\tau(x_0, x_1). \quad (5.3)$$

Let N be the normal subgroup generated by r in G^* . Our desired G will be G^*/N . Small cancellation theory provides us a way to decide when a word w in G^* belongs to N . This is expressed by Proposition 5.2. Any weakly cyclically reduced words in N must contain relatively “big parts” of words in the symmetrized closure R of $r(z, x_0, x_1)$. We need to recall the notions involved in this result. More details can be found in the book by Lyndon and Schupp [13, p. 286] or in Shelah [16, p. 378].

Let $g = g_1 g_2 \cdots g_m$ be a canonical representation of a word $g \in G^* = G_1 *_{G_0} G_2$. Let $|g| = m$ denote the *length* of g . Recall that g is *weakly cyclically reduced* if m is even, or $m = 1$ or $g_m g_1 \notin G_0$, equivalently, $g_1 \cdots g_m$ has no conjugate of length $< m - 1$. Note that any word is conjugate to a weakly cyclically reduced word. Let R be the *symmetrized closure* of $\{r(z, x_0, x_1)\}$ in G^* , i.e. add all inverses and all conjugates which are weakly cyclically reduced.

A *piece* (relative to R) is an element b in G^* for which there exist two different elements $r, r' \in R$ such that $r = bc$ and $r' = bc'$. Hence, a piece is a subword in R which cancels out in the multiplication of two non-inverse elements of R . We say that R satisfies the small cancellation condition $C'(\frac{1}{10})$ if *all pieces are relatively small*, that is, if for some $r \in R$, there is a piece b such that $r = bc$, then $|b| \leq \frac{1}{10}|r|$.

It is known from Lyndon and Schupp [13, p. 289] that if R is the symmetrized subset generated by the word $r(z, x_0, x_1)$ of (5.3), then R satisfies $C'(\frac{1}{10})$. In particular, Theorem 1.9 in Shelah [16] yields part 1 of the following proposition. Part 2 is obvious.

Proposition 5.2 *Let $G^* = G_1 *_{G_0} G_2$, R and N be as before. Let w be a weakly cyclically reduced word of $G^* \setminus \{1\}$. Then the following holds:*

1. *If $w \in N$, then w contains more than $\frac{7}{10}$ of an element r of R , that is, w has a subword w_1 which is also contained in r and such that $|w_1| \geq \frac{7}{10}|r|$.*
2. *If $w \notin N$, then w does not contain $\frac{1}{2}$ of any element $r \in R$, that is, no $r \in R$ has a subword w_1 also contained in w with $|w_1| \geq \frac{1}{2}|r|$. $\square$*

Proposition 5.3 *Let $G^* = G_1 *_{G_0} G_2$, R and N be as before. Let $G = G^*/N$. Then $G \in \mathcal{K}$, $G_0 \leq G$ and satisfies the properties (1), (2) and (3) stated before.*

PROOF. Since elements of G_1 and G_2 have length 1 in G^* , we have that G_1 and G_2 embed in $G = G^*/N$, by Proposition 5.2. It also follows that G_1 and G_2 are malnormal

subgroups of G ; see Lyndon and Schupp [13, p. 203] or Shelah [16, p. 383]. Hence G satisfies properties (1) and (2).

Let us prove that G is in $\mathcal{K}$: Let $H' \leq G$ be a subgroup isomorphic to H and $x \in G$ such that $[H', x] = 1$. We now apply a version of the Torsion Theorem, see [13, p. 288, Theorem 11.2 and remarks after the theorem] for a fixed element $h \neq 1$ (of finite order) in H' . Hence, there exists an element $g \in G$ such that the conjugate h^g belongs to G_1 or G_2 ; we may assume $h^g \in G_1$. From $[H', x] = 1$ follows $[h, x] = 1$, hence $[h^g, x^g] = 1$ and

$$1 \neq (h^g)^{x^g} = h^g \in G_1 \cap G_1^{x^g}.$$

Malnormality of G_1 in G implies that x^g also belongs to G_1 . By hypothesis $G_1 \in \mathcal{K}$, hence $x^g = 1$ and thus $x = 1$.

Finally we show that (3) holds. Let $\psi_0 : G_0 \rightarrow G_0$ be a given monomorphism which extends to a monomorphism ψ with domain G . Suppose that $\psi \upharpoonright H_0$ is conjugation by some element $y_0 \in G_1 \setminus G_0$. We shall assume for simplicity that $y_1 = 1$ (a similar argument will prove the general case). Since ψ extends ψ_0 we have

$$\tau(\psi(x_0), \psi(x_1)) = \psi(z) = \psi_0(z) = g_0$$

for some $g_0 \in G_0$. In other words, we have the following relation in G

$$g_0^{-1}(x_0^{y_0} x_1)(x_0^{y_0} x_1^2)(x_0^{y_0} x_1)^2(x_0^{y_0} x_1^2)^2 \cdots (x_0^{y_0} x_1)^{80}(x_0^{y_0} x_1^2)^{80} = 1.$$

Note that y_0 has length 1 in G^* and that $x_0^{y_0}$ cannot cancel with x_1 in G^* . Hence, if we view $g_0^{-1}y_0^{-1}$ as a new element, then the new word is written in a weakly cyclically reduced form in G^* . Proposition 5.2 tells us that this word must contain $\frac{7}{10}$ of some word of R . This forces $x_0^{y_0} = x_0$, that is, $y_0 \in C_{G_1}(x_0)$. But this contradicts the assumption made on y_0 . Hence $y_0 \in G_0$ as desired. $\square$

6 A universe for our group constructions. A uniform partially ordered set of approximations


Let λ be as in Main Theorem. We shall need in Lemma 6.7 below to assume GCH. Let H be a suitable group with automorphism group $\widehat{H}$ of size $< \lambda$.

First of all we need to define a “universe” where we can embed our groups. We also want to be sure that there is “enough room” for new elements which are added during the construction of our desired group. This suitable universe is the set

$$\widetilde{\lambda}^+ := \{\lambda\alpha + i : \alpha \in \lambda^+, i < \lambda\}. \quad (6.1)$$

Recall that this is the well-ordered set $\lambda^+ \times \lambda$ of cardinality λ^+ . If $x = \lambda\alpha + i \in \widetilde{\lambda}^+$ with $i < \lambda$, we write $\|x\| = \alpha$ and call α the *norm* of x . If X is a subset of $\widetilde{\lambda}^+$, the *domain* of X , which we denote by $\text{dom } X$, is the collection of all $\|x\|$ for all $x \in X$; and its norm $\|X\| := \|\sup X\|$ is the norm of the supremum $\sup X \in \widetilde{\lambda}^+$. This is well-defined if $|X| \leq \lambda$.

The following picture illustrates how we can embed for instance $H * H_\alpha$ in $\widetilde{\lambda}^+$, with $\text{dom}(H * H_\alpha) = \{0, \alpha\}$. More details are given in Proposition 6.4 below.



Definition 6.1 Let $u \subset \lambda^+$ be a subset of cardinality $< \lambda$ with $0 \in u$. Then a group G of size $|G| < \lambda$ is called an u -group if the following holds:

- (a) $\text{Dom } G \subset \widetilde{u} := \{\lambda\alpha + i : \alpha \in u, i < \lambda\}$, where $\text{Dom } G$ denotes the underlying set of elements of the group G . We often identify $\text{Dom } G = G$.
- (b) For every $\delta \in \widetilde{\lambda}^+$ divisible by λ , the subset $G \cap \delta$ is a subgroup of G .
- (c) $\text{dom } G = u$, that is, for every ordinal $\alpha \in u$, there exists an element $g \in G$ such that $\lambda\alpha \leq g < \lambda(\alpha + 1)$.

We now combine Definition 6.1 with the constructions given in Section 4. Consider two new families of groups $s\mathcal{K}$ and $s\mathcal{K}^*$:

Definition 6.2 1. A group G belongs to $s\mathcal{K}$ if G is an u -group for some $u \subset \lambda^+$, such that, for every δ divisible by λ , the subgroup $G \cap \delta$ belongs to $\mathcal{K}$ and satisfies the following condition:

If two copies of H in $G \cap \delta$ are conjugate by some element $g \in G$, then $g \in G \cap \delta$.

2. A group G belongs to $s\mathcal{K}^*$ if G is a member of $s\mathcal{K}$ such that for every ordinal δ divisible by λ , the subgroup $G \cap \delta$ belongs to $\mathcal{K}^*$.

Note that $G \in \mathcal{K}$ (resp. $G \in \mathcal{K}^*$) follow respectively from $G \in s\mathcal{K}$ (resp. $G \in s\mathcal{K}^*$) and $|G| < \lambda$ because we can choose $\delta \in \widetilde{\lambda^+}$ divisible by λ with $G \cap \delta = G$.

Notation 6.3 We will not need the old families $\mathcal{K}$ and $\mathcal{K}^*$ anymore. So from now on we rename $s\mathcal{K}$ by $\mathcal{K}$ and $s\mathcal{K}^*$ by $\mathcal{K}^*$, respectively.

For a fixed $u \subset \lambda^+$ with $|u| < \lambda$ we denote by $\mathcal{K}_u$ (resp. $\mathcal{K}_u^*$) the set of all u -groups in the new $\mathcal{K}$ (resp. in $\mathcal{K}^*$).

Proposition 6.4 $\mathcal{K}_u^*$ is dense in $\mathcal{K}_u$ for any $u \subset \lambda^+$ with $|u| < \lambda$ and $0 \in u$.

PROOF. The proof is the same as that of Proposition 4.11, but we must indicate how to assign ordinals conveniently to the constructions described in Section 4. In particular we next show that $\mathcal{K}_u$ is not empty. As $0 \in u$, we identify the automorphism group $\widehat{H}$ of H with the first $|\widehat{H}|$ ordinals in λ ; the identity element 1 goes to $0 \in \widetilde{\lambda^+}$. Similarly, for every $\alpha \in u$ we consider an isomorphic copy $\widehat{H}_\alpha$ of $\widehat{H}$, and we identify $\widehat{H}_\alpha \setminus \{1\}$ with the first $|\widehat{H}| - 1$ elements in $[\lambda\alpha, \lambda\alpha + \lambda)$. Let $G_0 = *_{\alpha \in u} \widehat{H}_\alpha$. Hence G_0 satisfies part (c) in Definition 6.1. If $x \in G_0$ then x has a unique reduced normal form $x = x_1 x_2 \cdots x_m$, where each x_i belongs to some factor $\widehat{H}_{\alpha_i}$ and successive factors are different. Let $\|x\|$ be the maximum of the norms $\|x_i\|$ for $i \leq m$. Now we assign to x any ordinal in $[\lambda\|x\|, \lambda\|x\| + \lambda)$. There is enough room in this segment, since λ is uncountable and $|H| \leq |\widehat{H}| < \lambda$, we simply choose any free place. Hence G_0 is an u -group. This yields a new version of Lemmas 4.2 and 4.3. The assignments for Lemma 4.9 are clear by continuity and note that we can do this because the length of the chains is strictly smaller than λ .

Suppose that G and G' have been already embedded in $\subset \widetilde{\lambda^+}$, and let $G^* = G *_{G_0} G'$ for some common subgroup G_0 . If $x \in G^*$ consider all possible reduced words $x = w(x_1, \dots, x_n)$ with $x_i \neq 1$ in one of the factors of G^* . Assign x to $\widetilde{\lambda^+}$ such that $\|x\|$ is the minimum of all norms of such words. The same idea works for the rest of results in Section 4. In particular, for HNN extensions we assign the new element t which conjugates $\widehat{H}$ and an isomorphic copy $\widehat{H}_1$ to a new ordinal $t \in \widetilde{u}$ such that $\|t\| = \|\widehat{H}_1\|$.

□

Lemma 6.5 $\mathcal{K}^*$ is closed under increasing chains of length $< \lambda$.

PROOF. The proof is straightforward. $\square$

Definition 6.6 Let G_i be u_i -groups for $i = 1, 2$. A group isomorphism $\psi : G_1 \rightarrow G_2$ is a *strong isomorphism* if $\text{Dom } \psi : \text{Dom } G_1 \rightarrow \text{Dom } G_2$ preserves the order in $\widetilde{\lambda}^+$ and it is constant modulo λ , i.e. for every $x \in \text{Dom } G_1$, $\psi(x) \equiv x \pmod{\lambda}$. This is equivalent to say that if $x = \lambda\alpha + i$ for some $\alpha \in u_1$ and $i < \lambda$, then $\psi(x) = \lambda\alpha' + i$ for some $\alpha' \in u_2$.

Clearly, the relation of being strongly isomorphic is an equivalence relation on $\mathcal{K}$ and hence on $\mathcal{K}^*$. Let $[\mathcal{K}]$ denote the set of equivalence classes $[G]$ for $G \in \mathcal{K}$ and $\pi : \mathcal{K} \rightarrow [\mathcal{K}]$ be the natural projection.

Lemma 6.7 (Assume GCH) For every $u \subset \lambda^+$ with $|u| < \lambda$ and $0 \in u$, the set $[\mathcal{K}_u]$ has cardinality $\leq \lambda$.

PROOF. For a fixed $u \subset \lambda^+$ with $|u| < \lambda$ we have exactly $\lambda^{<\lambda}$ different subsets in $\widetilde{u}$ of size $< \lambda$, and $\lambda^{<\lambda} = \lambda$ by GCH. Now the number of group structures supported by some set X is $\leq 2^{|X|}$. By GCH again and $|X| < \lambda$ we have $2^{|X|} = |X|^+ \leq \lambda$. We conclude $|[\mathcal{K}_u]| \leq \lambda$. $\square$

In fact $|[\mathcal{K}_u]| = \lambda$. This follows given other assignments in the proof of Proposition 6.4. It is based on a decomposition of u into λ subsets of size λ , which however will not be needed.

Hence there exists an inclusion (in fact a bijection) $\text{cod} : [\mathcal{K}_u] \rightarrow \lambda$, which induces a map $\text{cod} : \mathcal{K}_u \xrightarrow{\pi} [\mathcal{K}_u] \rightarrow \lambda$. We can assume that this bijection respects partial orders, i.e. if $G \leq G'$ then $\text{cod } G \leq \text{cod } G'$.

We finally collect all this information in a *coding* map

$$c : \mathcal{K} \rightarrow \lambda \times \mathcal{P}_{<\lambda}(\lambda^+), \quad \text{given by } G \mapsto (\text{cod } G, \text{dom } G),$$

with $c(G) = c(G')$ if G and G' are strongly isomorphic. Let $\mathbb{P} := c(\mathcal{K}^*)$. Then the partial order on $\mathcal{K}$ induces a partial order on $c(\mathcal{K})$ and in particular on $\mathbb{P}$ as follows: we say that $p \leq q$ if there are groups G_p and G_q in $\mathcal{K}$ coded by p and q , respectively, such that $G_p \leq G_q$. Observe that every element $p = (\alpha, u)$ in $\mathbb{P}$ codes every strong isomorphism class $[G]$ in $\mathcal{K}^*$, hence it is natural to view the elements of $\mathbb{P}$ as groups and to consider $\text{Aut}(p)$ for instance.

Theorem 6.8 $(\mathbb{P}, \leq)$ is a λ^+ -uniform partially ordered set.

PROOF. We have to check the conditions listed in Definition A.1.

1. Let $p, q \in \mathbb{P}$ such that $p \leq q$. Then there are groups $G_p, G_q \in \mathcal{K}^*$ coded by p and q respectively, such that G_p is a subgroup of G_q . This yields $\text{dom } G_p \subseteq \text{dom } G_q$, or equivalently $\text{dom } p \subseteq \text{dom } q$, as desired.

2. Let $p, q, r \in \mathbb{P}$ such that $p, q \leq r$. Let G_p, G_q and G_r groups in $\mathcal{K}^*$ coded by p, q and r , respectively, such that $G_p, G_q \leq G_r$. Consider G' the subgroup of G_r generated by G_p and G_q . It is clear that $\text{dom } G' = \text{dom } p \cup \text{dom } q$ and moreover $G' \in \mathcal{K}$. Using the fact that G_r is already in $\mathcal{K}^*$, we can add to G' the elements of G_r , with norm in the domain of G' , in order to obtain a new group $G'' \in \mathcal{K}^*$ such that $G' \leq G'' \leq G_r$. Finally, define $r' = c(G'')$.

The proofs of parts 3 to 6 are straightforward.

7. (Indiscernibility) Suppose that $p = (\alpha, u) \in \mathbb{P}$ and $\varphi : u \rightarrow u'$ is an order-isomorphism in λ^+ . Let $G_p \in \mathcal{K}^*$ be a group coded by p . Using φ we can define a strong isomorphism $\psi : G_p \rightarrow \psi(G) \subset \widetilde{\lambda^+}$ as follows: if $x = \lambda\zeta + i \in G_p$ for some $\zeta \in u \setminus \{0\}$ and $i < \lambda$, define $\psi(x) = \lambda\varphi(\zeta) + i$. Clearly, the image $\psi(G)$ is a group in $\mathcal{K}^*$ which has code $(\alpha, u') = \varphi(p)$. Hence $\varphi(p) \in \mathbb{P}$. The fact that $q \leq p$ implies $\varphi(q) \leq \varphi(p)$ is also clear.

8. (Amalgamation property) Let G_p and $G_q \in \mathcal{K}^*$ coded by p and q respectively. Let $G_0 \in \mathcal{K}^*$ coded by $p \restriction \alpha$, such that $G_0 \leq G_p$ and $G_0 \leq G_q$. Consider G^* the free product of G_p and G_q amalgamating the common subgroup G_0 . Then $G^* \in \mathcal{K}$ by Lemma 4.13. Now by Proposition 6.4 there is a group $G \in \mathcal{K}^*$ such that $G^* \leq G$. Take $r = c(G)$.


□

7 Small cancellation groups in $\widetilde{\lambda^+}$

In this section we formulate a refined version of Proposition 5.3. In the proof of our Main Theorem we will have to show that the final group $\mathcal{G}$ is complete and co-hopfian. We will suppose that a monomorphism $\psi : \mathcal{G} \rightarrow \mathcal{G}$ exists which is not an inner automorphism. Using the help of some prediction principles, we will be able to find restrictions of ψ on many layers of our group $\mathcal{G}$. In Proposition 7.1 we provide a method to get rid of such monomorphism ψ .

Recall that $\mathcal{K}$ and $\mathcal{K}^*$ are the new families introduced in Notation 6.3.

Proposition 7.1 *Let $\delta_0 < \delta_1 < \delta_2$ be ordinals in $\widetilde{\lambda}^+$ of cofinality λ . Suppose that we have G_i, ψ_i, H_i, y_i, z as in the following picture*



where

1. $G_i \in \mathcal{K}^*$ for $i = 0, 1, 2$.
2. $G_i \subseteq \delta_i$ and $G_i \cap \delta_0 = G_0$ for $i = 1, 2$.
3. $\psi_i : G_i \rightarrow G_i$ are not inner automorphisms such that $\psi_i \upharpoonright G_0 = \psi_0$ for $i = 1, 2$.
4. $H_0 \leq G_1$ and $H_1 \leq G_2$ are two copies of H such that $H_i \cap \delta_i = \{1\}$ for $i = 0, 1$.
5. $\psi_i \upharpoonright H_i = y_i^* \upharpoonright H_i$ for some $y_i \in G_{i+1}$ for $i = 0, 1$.
6. $z \in G_0$.

Choose $1 \neq x_i \in H_i$ such that ψ_i is not conjugation by an element in $C_{G_{i+1}}(x_i)$ for $i = 0, 1$ and let $\tau(x_0, x_1)$ be as in (5.2). Then there is a small cancellation group G in $\mathcal{K}$ satisfying the following:

- (a) The relation $z = \tau(x_0, x_1)$ holds in G .
- (b) G_1 and G_2 are malnormal subgroups of G .
- (c) If ψ is an endomorphism of G extending ψ_1 and ψ_2 and for some $i = 0, 1$, $\psi \upharpoonright H_i$ is conjugation by some element $y_i \notin G_0$, then $\tau(\psi(x_0), \psi(x_1)) \notin G_0$.

PROOF. Using Proposition 5.3, the desired group is the free group $G^* = G_1 *_{G_0} G_2$ modulo the small cancellation $z = \tau(x_0, x_1)$. More precisely G is the quotient of G^* by the normal subgroup N generated by the symmetrized closure R of $r = z^{-1}\tau(x_0, x_1)$. Hence (a), (b) and (c) hold. $\square$

8 Proof of the Main Theorem

In this section we finally prove the existence of the group $\mathcal{G}$ satisfying the properties listed in the Main Theorem.

Theorem A.4 will ensure that under Dl_λ player II has a winning strategy for the $\mathbb{P}$ -game, where $\mathbb{P}$ is the λ^+ -uniform partially ordered set constructed in Section 6. In other words player II will be able to find an increasing and unbounded sequence of ordinals ζ_α in λ^+ and an ascending chain of admissible ideals $\mathbb{G}_\alpha$ on $\mathbb{P}_{\zeta_\alpha}$ which meet the density systems defined by player I. In the following Subsections 8.1, 8.2 and 8.3 we will tell player I how he must define these density systems. These restrictions will guarantee that our final group $\mathcal{G}$ has cardinality λ^+ , is complete (d), co-hopfian (e) and simple (f). Conditions (a), (b) and (c) of the Main Theorem will be automatically satisfied, since all the groups in the direct system are in $\mathcal{K}^*$.

The group $\mathcal{G}$ at the end will be

$$\mathcal{G} = \bigcup_{\alpha < \lambda^+} \mathcal{G}_\alpha, \quad (8.1)$$

where each $\mathcal{G}_\alpha$ is the union of all groups (coded) in the directed system $\mathbb{G}_\alpha$. Every $\mathcal{G}_\alpha$ has cardinality $\leq \lambda$, so that (8.1) is a λ^+ -filtration of $\mathcal{G}$.

Let us start playing the $\mathbb{P}$ -game. We proceed by transfinite induction on $\alpha < \lambda^+$. By continuity we shall only consider the induction step from α to $\alpha + 1$. Suppose that player II has already found an ordinal $\zeta_\alpha < \lambda^+$ and an admissible ideal $\mathbb{G}_\alpha$ in $\mathbb{P}_{\zeta_\alpha}$ where

$$\mathbb{P}_{\zeta_\alpha} = \{p \in \mathbb{P} : \text{dom } p \subseteq \zeta_\alpha\}.$$

Now it is the turn of player I. He chooses $p_{\alpha+1} \in \mathbb{P}/\mathbb{G}_\alpha = \{p \in \mathbb{P} : p \restriction \zeta_\alpha \in \mathbb{G}_\alpha\}$, such that $\|p_{\alpha+1}\| > \zeta_\alpha$. This will guarantee that the length of the game is λ^+ . He can also define, for each $\zeta_\alpha < \beta < \lambda^+$, a list of (ζ_α, β) -density systems D^i over $\mathbb{G}_\alpha$, where i ranges over a set of cardinality $\leq \lambda$.

Given $u \subseteq v \subseteq \lambda^+$ with $|v| < \lambda$, we denote for the rest of this section

$$\mathbb{E} := \{p \in \mathbb{P}/\mathbb{G}_\alpha : \text{dom } p \subseteq v \cup \beta\}.$$

Next we define λ subsets $D(u, v)$ in $\mathbb{E}$ satisfying the properties stated in Definition A.3. In particular, property 2 follows directly if we define in the same way $D(u', v')$ for all pairs (u', v') such that $u \cap \beta = u' \cap \beta$, $v \cap \beta = v' \cap \beta$ and there is an order-isomorphism from v onto v' which maps u onto u' .

8.1 Density system to obtain $|\mathcal{G}| = \lambda^+$

We want that our group $\mathcal{G}$ has $\text{dom } \mathcal{G} = \lambda^+$. This will imply that $|\mathcal{G}| = \lambda^+$. With this objective in mind player I chooses in his α th move the following density system:

If u, v are subsets in λ^+ such that $u \subseteq v$ and $|v| < \lambda$ player I defines

$$D(u, v) := \{p \in \mathbb{E} : \text{if } \alpha \notin u \text{ then } \alpha \in \text{dom } p\}.$$

Proposition 8.1 *D is a density system over $\mathbb{G}_\alpha$.*

PROOF. It is clear that $D(u, v)$ is closed upwards in $\mathbb{E}$. Now we have to show that $D(u, v)$ is dense in $\mathbb{E}$. In the case when $\alpha \in u$ then $D(u, v) = \mathbb{E}$ and density is obvious. Suppose that $\alpha \notin u$ and let q be any element in $\mathbb{E}$. Let G be an u -group which is coded by q . As in the proof of Proposition 6.4 we can find a group $G' \in \mathcal{K}^*$, such that $G \leq G'$, which contains a subgroup H' isomorphic to H and $\alpha \in \text{dom } H' \subseteq [\lambda\alpha, \lambda\alpha + \lambda)$. In particular, we have $\alpha \in \text{dom } G'$. Moreover, the group G' can be constructed in such a way that $\text{dom } G' \subseteq v \cup \beta$. Therefore, $p := (\text{cod } G', \text{dom } G') \in D(u, v)$ and $q \leq p$ as desired. $\square$

8.2 Density systems to obtain that $\mathcal{G}$ is complete and co-hopfian

Suppose for a moment that $\mathcal{G}$ is already constructed as a limit of groups in $\mathcal{K}^*$ and ψ is an unwanted monomorphism of $\mathcal{G}$ which is not a inner automorphism. Hence, it is not inner on many layers by Proposition 5.1.

First we note that for a fixed pair (u, v) and $\zeta_\alpha < \beta < \lambda^+$ the set E of equivalence classes of all monomorphisms ‘up to order isomorphism’ of u -groups U with $u \subseteq v \cup \beta$ has size λ . Choose then an enumeration of $U \times E$ by λ . For every $i \in \lambda$ we shall define a (ζ_α, β) -density system D^i taking care of an element $z \in U$ and the monomorphism ψ coded by $i \in \lambda$. If $p \in \mathbb{E}$ then consider $p \restriction u = G_0$ and let $z \in G_0, \psi \in \text{Aut}(G_0)$ be the pair of elements coded by $i < \lambda$. Then we put $p \in D^i(u, v)$ if ψ is not inner and G_0 extends to $p = G$ with $G_1, G_2, \psi_1, \psi_2, H_0$ and H_1 as in Proposition 7.1.

Note that such a monomorphism of G_0 can not survive to a monomorphism of G . On the other hand if $p \in \mathbb{E}$ such that ψ is inner on $p \restriction u$ then $p \notin D^i(u, v)$.

Proposition 8.2 D^i is a density system over $\mathbb{G}_\alpha$ for every $i \in \lambda$.

PROOF. It is clear that every $D^i(u, v)$ is closed upwards. The fact that $D^i(u, v)$ is dense in $\mathbb{E}$ is shown in Proposition 7.1, where we can find a small cancellation group $G \in \mathcal{K}$ with the required properties. If necessary, use also Proposition 6.4 to obtain a group G_3 in $\mathcal{K}^*$ such that $G \leq G_3$. $\square$

8.3 Density systems to obtain that $\mathcal{G}$ is simple

Finally we take care of simplicity of our group $\mathcal{G}$. As in the previous subsection, we shall define λ density systems D^i . As before, let $u \subseteq v \subseteq \lambda^+$ such that $|v| < \lambda$. Suppose that U is a group in $\mathcal{K}^*$ with $\text{dom } U = u$. We consider the set P_U of all ordered pairs (x, y) where $1 \neq y, x \in U$ and choose a coding map $c : P_U \hookrightarrow \lambda$. Then, for every $i \in c(P_U)$ player I chooses $D^i(u, v)$ to be

$$\{p \in \mathbb{E} : p = c(G), p \upharpoonright u = c(U), c(x, y) = i, \text{ and } x \notin \langle y^U \rangle \implies x \in \langle y^G \rangle\},$$

where $\langle y^U \rangle = \langle y^z : z \in U \rangle$.

Proposition 8.3 D^i is a density system over $\mathbb{G}_\alpha$ for every i .

PROOF. Let $q \in \mathbb{E}$ represent the group U and let $1 \neq y, x \in U$ be such that $c(x, y) = i$. If $x \in \langle y^U \rangle$, then $D^i(u, v) = \mathbb{E}$ and density is obvious. Suppose that $x \notin \langle y^U \rangle$. We have to construct a v -group G represented by $p \in \mathbb{E}$ extending q such that $x \in \langle y^G \rangle$.

We consider the three possible cases:

Case 1: $o(x) = o(y) = \infty$. In this case we use Proposition 4.14 as follows. Define $G_1 = \langle U, t : x = t^{-1}yt \rangle$, where t is a new element in v with $||t|| = ||x||$. Of course, $G_1 \in \mathcal{K}$, so we can construct $G_1 \leq G$ with $G \in \mathcal{K}^*$ and $\text{dom}(G) \subset v$, as explained in the proof of Proposition 6.4. Hence x and y are conjugate in G .

Case 2: $o(x) = n < \infty$ and $o(y) = \infty$. Here, we use Lemma 4.8 (part 2 of its proof). We can find an element $x' = xz \in G \in \mathcal{K}^*$ of infinite order such that $z^{-1}y$ has infinite order as well. From case 1 we note that y, xz and z^{-1} are conjugate in G . Hence $x \in \langle y^G \rangle$.

Case 3: If y has finite order, we argue similarly. Take a conjugate z of y in some $G \in \mathcal{K}^*$ such that yz has infinite order. Hence $\langle y^G \rangle$ contains all elements of infinite order and case 1 and 2 apply.

We conclude in any case that D^i is a density system. $\square$

8.4 Existence of $\mathcal{G}$

Using the density systems and the winning strategy of the game Theorem A.4 we find the building blocks for the desired group.

Theorem 8.4 (Assume DI_λ) *There is a chain of admissible ideals $\mathbb{G}_\alpha \subseteq \mathbb{P}_{\zeta_\alpha}$ for an unbounded sequence $\{\zeta_\alpha < \lambda^+ : \alpha < \lambda^+\}$ such that $\mathbb{G} = \bigcup_{\alpha < \lambda^+} \mathbb{G}_\alpha$ satisfies the following conditions.*

1. *For every $\alpha \in \lambda^+$ there is a group $G \in \mathbb{G}$ such that $\alpha \in \text{dom } G$.*
2. *Completeness and co-hopfian: If $\delta_0 < \widetilde{\lambda}^+$ divisible by λ , $z \in G_0 \in \mathbb{G}$ with $G_0 \subseteq \delta_0$ and $\psi_0 : G_0 \rightarrow G_0$ is monomorphism which is not inner, then there exist ordinals $\delta_0 < \delta_1 < \delta_2$ in $\widetilde{\lambda}^+$ divisible by λ and groups $G_0 \leq G_i \in \mathbb{G}$ with $G_i \subseteq \delta_i$ and subgroups $H_0 \leq G_1$, $H_1 \leq G_2$ isomorphic to H such that $H_i \cap \delta_i = \{1\}$ for $i = 0, 1$ as in the hypothesis of Proposition 7.1.*
3. *Simplicity: For every $1 \neq y, x \in G \in \mathbb{G}$ there is $G' \in \mathbb{G}$ such that $G \leq G'$ and $x \in \langle y^{G'} \rangle$.*

PROOF. We apply Theorem A.4 and let player I choose the density systems above. Condition 1 of the Theorem follows immediately from Proposition 8.1, condition 2 follows from Proposition 8.2 and condition 3 follows from Proposition 8.3. $\square$

Let $\mathcal{G}$ be the group which is the union of the directed system $\mathbb{G}$ given in Theorem 8.4. We finally check that $\mathcal{G}$ has the desired properties:

$\mathcal{G}$ has cardinality λ^+ : By Theorem 8.4 we have that for every $\alpha < \lambda^+$, there is a group G in $\mathbb{G}$ such that $\alpha \in \text{dom } G$. This says that $\text{dom } \mathcal{G} = \lambda^+$ and hence, $\lambda^+ \leq |\mathcal{G}|$. On the other hand, all groups $G \in \mathbb{G}$ lie in $\widetilde{\lambda}^+$. Thus $|\mathcal{G}| \leq |\widetilde{\lambda}^+| = \lambda^+$ and $|\mathcal{G}| = \lambda^+$ follows.

Properties (a)-(c): These properties are satisfied by all the groups G in $\mathbb{G}$ by construction. Now if $x \in \mathcal{G}$ and $H_1 \leq \mathcal{G}$ is isomorphic to H such that $[H_1, x] = 1$ there exists a group $G \in \mathbb{G}$ containing x and H_1 . Hence $x = 1$ from $G \in \mathcal{K}^*$. This shows property (c). The properties (a) and (b) can be proved similarly.

Properties (d)-(e): Let $\psi : \mathcal{G} \rightarrow \mathcal{G}$ be a monomorphism (possibly not inner automorphism). Let C be the set of ordinals $\delta < \widetilde{\lambda}^+$ which are divisible by λ and such that ψ maps $\delta \cap \mathcal{G}$ into $\delta \cap \mathcal{G}$. Then C is clearly a club of $\widetilde{\lambda}^+$. Now let $S = \{\delta \in \widetilde{\lambda}^+ : \text{cf}(\delta) = \lambda\}$. Then S is stationary in $\widetilde{\lambda}^+$ and hence $S \cap C \neq \emptyset$.

Choose for each $\delta \in S$ a subgroup H_δ^0 of $\mathcal{G}$ isomorphic to H such that

$$H_\delta^0 \cap \delta = \{1\}.$$

By property (b) we find $y_\delta \in \mathcal{G}$ such that $\psi \upharpoonright H_\delta^0 = y_\delta^* \upharpoonright H_\delta^0$.

Now we have two cases:

Case 1: Suppose that $y_\delta < \delta$ for all $\delta \in S$. In that case we have a regressive function

$$S \rightarrow \widetilde{\lambda^+}, \quad \text{given by } \delta \mapsto y_\delta.$$

Hence, by Fodor's lemma (see Lemma 3.1), there exists a stationary subset $S' \subset S$ and $y \in \mathcal{G}$ such that $y_\delta = y$ for all $\delta \in S'$.

Let $z \in \mathcal{G}$. We shall prove that $\psi(z) = y^*(z)$. For every $\alpha \in \lambda^+$ choose an ordinal $\beta = \beta(\alpha) > \alpha$ such that $\lambda\beta \in S'$. Such a β exists since S' is stationary and $\{\beta \in \lambda^+ : \beta \geq \alpha\}$ is a club. Let $G_\alpha \in \mathbb{G}$ such that $z \in G_\alpha$ and $H_\alpha = H_\beta^0 \leq G_\alpha$. Let $H'_\alpha = \psi(H_\alpha)$. Note that H'_α is a subgroup of G_α such that $(\text{dom } H'_\alpha) \cap \alpha = \emptyset$. Now by Theorem 8.4 we can find ordinals $\alpha_0 < \alpha_1 < \alpha_2$, groups G_0, G_1 and G_2 in $\mathbb{G}$ satisfying the hypothesis of Proposition 7.1 (take $\delta_i = \lambda\alpha_i$). Then there exists a group $G \in \mathbb{G}$ extending G_1 and G_2 where the small cancellation $z = \tau(x_0, x_1)$ holds, where $z \in G_0$ and $x_i \in H_i$ for $i = 0, 1$. Hence

$$\psi(z) = \psi(\tau(x_0, x_1)) = \tau(\psi(x_0), \psi(x_1)) = \tau(y^*(x_0), y^*(x_1)) = y^*\tau(x_0, x_1) = y^*(z).$$

Since z is arbitrary, we have shown that ψ is an inner automorphism.

Case 2: Suppose that for some $\delta \in S$, we have $\psi \upharpoonright H_\alpha^0 \neq y^* \upharpoonright H_\alpha^0$ for any $y \in \delta$. We argue as in the case 1 with $z = 1$. Hence, there exist ordinals $\alpha_0 < \alpha_1 < \alpha_2$ and groups $G_i \in \mathcal{K}^*$ as in Proposition 7.1 so that $1 = z = \tau(x_0, x_1)$ for some elements x_0 and x_1 . Hence $1 = \psi(z) = \tau(\psi(x_0), \psi(x_1)) \notin G_0$, which is impossible.

Thus $\mathcal{G}$ is complete and co-hopfian, as desired.

Property (f): This follows from the third condition of Theorem 8.4.

A The combinatorial principle Dl_λ

In this appendix we recall for convenience of the reader the combinatorial result proved by Hart, Laflamme and Shelah in [9]. This result will guarantee in Theorem 8.4 the existence of a certain direct system $\mathbb{G}$ and $\mathcal{G}$ in Main Theorem will be the limit of the groups coded by elements in $\mathbb{G}$.

Theorem A.4 is stated in terms of a game of λ^+ moves between two players I and II. In the turn $\alpha < \lambda^+$ player II picks an ordinal $\zeta_\alpha < \lambda^+$ and an admissible ideal $\mathbb{G}_\alpha$ from a partial uniform set $\mathbb{P}$ (see definitions below).

The elements of $\mathbb{P}$ are pairs $p = (\alpha, u) \in \lambda \times \mathcal{P}_{<\lambda}(\lambda^+)$. We write $u = \text{dom } p$ and call u the *domain* of p . The ordinals α are there to code all possible algebraic structures that u can support. In our case α will run over all strongly isomorphic classes of groups supported by a fixed u (see Section 6).

Definition A.1 A *standard λ^+ -uniform partial order* is a partial order $\leq$ defined on a subset $\mathbb{P}$ of $\lambda \times \mathcal{P}_{<\lambda}(\lambda^+)$ satisfying the following conditions:

1. (Compatibility of the orders) If $p \leq q$ then $\text{dom } p \subseteq \text{dom } q$.
2. For all $p, q, r \in \mathbb{P}$ with $p, q \leq r$ there is $r' \in \mathbb{P}$ such that $p, q \leq r' \leq r$ and $\text{dom } r' = \text{dom } p \cup \text{dom } q$.
3. If $\{p_\alpha : \alpha < \delta\}$ is an increasing sequence in $\mathbb{P}$ of length $\delta < \lambda$ then it has a least upper bound $q \in \mathbb{P}$ with $\text{dom } q = \bigcup_{\alpha < \delta} \text{dom } p_\alpha$; we say that $q = \bigcup_{\alpha < \delta} p_\alpha$.
4. If $p \in \mathbb{P}$ and $\alpha < \lambda^+$ then there is $q \in \mathbb{P}$ such that $q \leq p$ and $\text{dom } q = \text{dom } p \cap \alpha$ and there is a unique maximal such q for which we write $q = p \restriction \alpha$.
5. (Continuity I) If δ is a limit then $p \restriction \delta = \bigcup_{\alpha < \delta} p \restriction \alpha$.
6. (Continuity II) If $\{p_\alpha : \alpha < \delta\}$ is an increasing sequence in $\mathbb{P}$ of length $\delta < \alpha$ then

$$\left(\bigcup_{i < \delta} p_i\right) \restriction \alpha = \bigcup_{i < \delta} (p_i \restriction \alpha).$$

7. (Indiscernibility) If $p = (\alpha, u) \in \mathbb{P}$ and $\varphi : u \rightarrow u'$ is an order-isomorphism in λ^+ , then $\varphi(p) := (\alpha, \varphi(u)) \in \mathbb{P}$. Moreover, if $q \leq p$ then $\varphi(q) \leq \varphi(p)$.
8. (Amalgamation property) For every $p, q \in \mathbb{P}$ and $\alpha < \lambda^+$ with $p \restriction \alpha \leq q$ and $\text{dom } p \cap \text{dom } q = \text{dom } p \cap \alpha$ there is an $r \in \mathbb{P}$ such that $p, q \leq r$.

If all these conditions hold, $\mathbb{P} = (\mathbb{P}, \leq)$ is called a *λ^+ -uniform partially ordered set*.

Consider the following filtration of $\mathbb{P}$, where $\alpha < \lambda^+$,

$$\mathbb{P}_\alpha := \{p \in \mathbb{P} : \text{dom } p \subseteq \alpha\}.$$

Definition A.2 Let $(\mathbb{P}, \leq)$ be a λ^+ -uniform partially ordered set and $\alpha < \lambda^+$. A subset $\mathbb{G} \subseteq \mathbb{P}_\alpha$ is an *admissible ideal* of $\mathbb{P}_\alpha$ if the following holds:

1. $\mathbb{G}$ is closed downward, i.e. if $p \in \mathbb{G}$ and $q \leq p$ then $q \in \mathbb{G}$.
2. $\mathbb{G}$ is λ -directed, i.e. if $\mathcal{A} \subset \mathbb{G}$ and $|\mathcal{A}| < \lambda$ then $\mathcal{A}$ has an upper bound in $\mathbb{G}$.
3. (Maximality) If $p \in \mathbb{P}$ is compatible with all $q \in \mathbb{G}$ then $p \in \mathbb{G}$. (Recall that p is compatible with q if there is an $r \in \mathbb{P}$ such that $p, q \leq r$.)

For an admissible ideal $\mathbb{G}$ of $\mathbb{P}_\alpha$, we define

$$\mathbb{P}/\mathbb{G} := \{p \in \mathbb{P} : p \restriction \alpha \in \mathbb{G}\}.$$

This consists of all extensions of elements in $\mathbb{G}$. Note that this notion is compatible with taking direct limits in the following sense. Let $\{\zeta_\beta : \beta < \alpha\}$ be an increasing sequence of ordinals in λ^+ converging to ζ . Suppose that for every β we have an admissible ideal $\mathbb{G}_\beta$ of $\mathbb{P}_{\zeta_\beta}$. Then there is a unique minimal admissible ideal containing the set theoretic union of the $\mathbb{G}_\beta$'s. With a slight misuse of notation we write

$$\mathbb{G}_{<\alpha} := \bigcup_{\beta < \alpha} \mathbb{G}_\beta,$$

for this admissible ideal of $\mathbb{P}_\zeta$, see Hart, Laflamme and Shelah [9, p. 173, Lemma 1.3].

Definition A.3 Let $\mathbb{G}$ be an admissible ideal of $\mathbb{P}_\alpha$ and $\alpha < \beta < \lambda^+$. An (α, β) -*density system* over $\mathbb{G}$ is a function

$$D : \{(u, v) : u \subseteq v \in \mathcal{P}_{<\lambda}(\lambda^+)\} \longrightarrow \mathcal{P}(\mathbb{P})$$

such that the following holds:

1. $D(u, v) \subseteq \{p \in \mathbb{P}/\mathbb{G} : \text{dom } p \subseteq v \cup \beta\}$ is a dense and upward-closed subset.
2. If $(u, v), (u', v')$ and $u \cap \beta = u' \cap \beta$, $v \cap \beta = v' \cap \beta$ and there is an order-isomorphism from v onto v' which maps u onto u' , then for any ordinal γ we have

$$(\gamma, v) \in D(u, v) \iff (\gamma, v') \in D(u', v').$$

An admissible ideal $\mathbb{G}'$ of $\mathbb{P}_{\alpha'}$ for some $\alpha' < \lambda^+$ *meets* the (α, β) -density system D (over $\mathbb{G}$) if $\alpha < \alpha'$, $\mathbb{G} \subseteq \mathbb{G}'$ and for each $u \in \mathcal{P}_{<\lambda}(\alpha')$ there is a $v \in \mathcal{P}_{<\lambda}(\alpha')$, with $u \subseteq v$ and such that $D(u, v) \cap \mathbb{G}' \neq \emptyset$.

The $\mathbb{P}$ -game

1. Player I moves first.

Let α be a successor ordinal and suppose that player I has played his $(\alpha - 1)th$ move.

2. Then player II picks an ordinal $\zeta_\alpha < \lambda^+$ and an admissible ideal $\mathbb{G}_\alpha$ of $\mathbb{P}_{\zeta_\alpha}$.
3. Then player I chooses an element $p_{\alpha+1} \in \mathbb{P}/\mathbb{G}_\alpha$ and chooses, for every $\zeta_\alpha < \beta < \lambda^+$, a list of (ζ_α, β) -density systems D^i over $\mathbb{G}_\alpha$, where $i \in I_\alpha$ with $|I_\alpha| \leq \lambda$.

If $\alpha = \lim_{\alpha' < \alpha} \alpha'$ is a limit ordinal the moves go as follow:

4. At the αth move, player II will have chosen an increasing sequence of ordinals $\zeta_{\alpha'} < \lambda^+$ and will have defined an admissible ideal $\mathbb{G}_{\alpha'}$ of $\mathbb{P}_{\zeta_{\alpha'}}$, for all $\alpha' < \alpha$.
5. Then player I will choose an element $p_\alpha \in \mathbb{P}/\mathbb{G}_{<\alpha}$ and he will also choose at most λ density systems D^i over $\mathbb{G}_{<\alpha}$, where $i \in I_\alpha$ with $|I_\alpha| \leq \lambda$.
6. After player I has finished his αth move player II will pick again an ordinal ζ_α and an admissible ideal $\mathbb{G}_\alpha$ of $\mathbb{P}_{\zeta_\alpha}$.

Player II *has a winning strategy* for the $\mathbb{P}$ -game if the sequences ζ_α and $\mathbb{G}_\alpha$ are increasing, we have $p_\alpha \in \mathbb{G}_\alpha$ and for all $\beta \geq \alpha$, $\mathbb{G}_\beta$ *meets* D_α^i for all $i \in I_\alpha$.

Theorem A.4 below states that player II has a winning strategy if Dl_λ holds. This is a combinatorial principle which helps player II to predict the moves of its opponent player.

Dl_λ asserts that there are sets $\mathcal{A}_\alpha \subseteq \mathcal{P}(\alpha)$ for all $\alpha < \lambda$, with $|\mathcal{A}_\alpha| < \lambda$ such that for all $X \subseteq \lambda$ the set $\{\alpha \in \lambda : X \cap \alpha \in \mathcal{A}_\alpha\}$ is stationary in λ .

Theorem A.4 *If Dl_λ holds then player II has a winning strategy for the $\mathbb{P}$ -game.*

We finally remark that $\lambda = \aleph_0$ or $\diamond(\lambda)$ implies Dl_λ . Recall that $\diamond(\lambda)$ is the particular case of Dl_λ where each $\mathcal{A}_\alpha$ consists on one single subset of λ . Moreover, if $\lambda = \kappa^+$ with $\text{cf}(\kappa) > \aleph_0$ and GCH holds, then Dl_λ holds. We also note that Dl_λ implies $\lambda^{<\lambda} = \lambda$.

References

- [1] M. Aubry, *Homotopy theory and models, Based on lectures held at a DMV Seminar in Blaubeuren by H. J. Baues, S. Halperin and J.-M. Lemaire*, DMV Seminar, Vol. **24**, Birkhäuser Verlag, Basel, 1995.
- [2] C. Casacuberta, *Recent advances in unstable localization*, CRM Proceedings and Lecture Notes **6**, Amer. Math. Soc., Providence, 1994, 1–22.
- [3] C. Casacuberta, J. L. Rodríguez and J.-Y. Tai, *Localizations of abelian Eilenberg–Mac Lane spaces of finite type*, preprint 1998.
- [4] C. Casacuberta, D. Scevenels and J. H. Smith, *Implications of large-cardinal principles in homotopical localization*, preprint 1998.
- [5] E. Dror Farjoun, *Cellular spaces, null spaces and homotopy localization*, Lecture Notes in Math. vol. 1622, Springer-Verlag, Berlin–Heidelberg–New York 1996.
- [6] M. Dugas, R. Göbel, *On locally finite p -groups and a problem of Philip Hall’s*, J. Algebra **159**(1) (1993), 115–138.
- [7] M. Dugas, A. Mader and C. Vinsonhaler, *Large E -rings exist*, J. Algebra **108** (1987), 88–101.
- [8] P. C. Eklof and A. H. Mekler, *Almost Free Modules. Set-theoretic Methods*, North-Holland, Amsterdam, New York 1990.
- [9] B. Hart, C. Laflamme and S. Shelah, *Models with second order properties V: A general principle*, Ann. Pure Appl. Logic, **64** (1993), 169–194.
- [10] T. Jech, *Set Theory*, Academic Press, New York, 1978.
- [11] A. Libman, *Cardinality and nilpotency of localizations of groups and G -modules*, Israel J. Math., to appear.
- [12] A. Libman, *A note on the localization of finite groups*, preprint 1998.
- [13] R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*, Springer Ergebnisseberichte **89**, Berlin–Heidelberg–New York 1977.

- [14] D. Robinson, *A Course in the Theory of Groups*, Graduate Texts in Math. vol **80**, Berlin–Heidelberg–New York 1996.
- [15] P. E. Schupp, *A characterization of inner automorphisms*, Proc. Amer. Math. Soc. **101** (1987), 226–228.
- [16] S. Shelah, *On a problem of Kurosh, Jonsson groups and applications*, S. I. Adian, W. W. Boone, G. Higman, eds., Word Problems II, North-Holland, Amsterdam, 1980, 373–394.
- [17] L. Strüngmann, *Almost-free $E(R)$ -algebras over countable domains*, Ph. D. Thesis, University of Essen (1998).
- [18] G. W. Whitehead, *Elements of Homotopy Theory*, Graduate Texts in Math. vol **61**, Springer-Verlag, Berlin–Heidelberg–New York, 1978.

Fachbereich 6, Mathematik und Informatik, Universität Essen,
D-45117 Essen, Germany, e-mail: `R.Goebel@Uni-Essen.De`

Departament de Matemàtiques, Universitat Autònoma de Barcelona,
E-08193 Bellaterra, Spain, e-mail: `jlrodri@mat.uab.es`

Institute of Mathematics, Hebrew University,
91904 Jerusalem, Israel, e-mail: `shelah@math.huji.ac.il`